



US Army Corps
of Engineers
Waterways Experiment
Station

Wetlands Research Program Technical Report WRP-SM-15

Evaluation of Mechanical and Chemical Methods for Control of *Melaleuca quinquenervia* in Southern Florida

by Alfred F. Cofrancesco, Jr., Jean W. Wooten, Harvey L. Jones

19960124 108



The following two letters used as part of the number designating technical reports of research published under the Wetlands Research Program identify the area under which the report was prepared:

	<u>Task</u>		<u>Task</u>
CP	Critical Processes	RE	Restoration & Establishment
DE	Delineation & Evaluation	SM	Stewardship & Management

The contents of this report are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such commercial products.

Evaluation of Mechanical and Chemical Methods for Control of *Melaleuca quinquenervia* in Southern Florida

by Alfred F. Cofrancesco, Jr., Harvey L. Jones

U.S. Army Corps of Engineers
Waterways Experiment Station
3909 Halls Ferry Road
Vicksburg, MS 39180-6199

Jean W. Wooten

University of Southern Mississippi
P.O. Box 5018
Hattiesburg, MS 39401


Final report

Approved for public release; distribution is unlimited

Prepared for **U.S. Army Corps of Engineers**
Washington, DC 20314-1000



**US Army Corps
of Engineers**
Waterways Experiment
Station



Waterways Experiment Station Cataloging-in-Publication Data

Cofrancesco, Alfred F.

Evaluation of mechanical and chemical methods for control of *Melaleuca quinquenervia* in southern Florida / by Alfred F. Cofrancesco, Jr., Harvey L. Jones, Jean W. Wooten ; prepared for U.S. Army Corps of Engineers.

34 p. : ill. ; 28 cm. -- (Technical report ; WRP-SM-15) (Wetlands Research Program technical report ; WRP-SM-15)

Includes bibliographic references.

1. *Melaleuca quinquenervia* -- Florida. 2. Trees -- Growth. 3. Herbicides. I. Jones, Harvey L. II. Wooten, Jean W. III. United States. Army. Corps of Engineers. IV. U.S. Army Engineer Waterways Experiment Station. V. Wetlands Research Program (U.S.) VI. Title. VII. Series: Wetlands Research Program technical report ; WRP-SM-15. VIII. Series: Technical report (U.S. Army Engineer Waterways Experiment Station) ; WRP-SM-15. TA7 W34 no. WRP-SM-15



Wetland Pests

Evaluation of Mechanical and Chemical Methods for Control of Melaleuca quinquenervia in Southern Florida (TR WRP-SM-15)

ISSUE:

Melaleuca quinquenervia (Cav.) Blake, common names melaleuca, cajeput, or punk tree, is native to the Australian region. Trees were first planted in southern Florida early in the 20th century. Starting in the late 1940s, melaleuca was planted by U.S. Army Engineer personnel at Lake Okeechobee, Florida, to serve as wind barriers and to provide protection for the levees surrounding the lake. The trees have now spread to wetland habitats and were found by a 1979 survey to grow along 88.5 km (53.1 miles) of the 175 km (105 miles) of levee road and to cover approximately 96.8 ha (242 acres) in these perimeter stands (Stocker 1982a). Additionally, Stocker found that "in the lake proper, melaleuca is found in approximately 420 ha (1,050 acres), where it offers as isolated individuals and as dense colonies (heads) up to 0.6 ha (1.5 acres) in size." This distribution of trees was approximately 0.3 percent of the lake acreage.

RESEARCH:

The objectives of this study were to determine the effects of selected treatment methods on vegetation regrowth and to assess their results in control of melaleuca. Seven approximately 3.3-ha plots were established in 1990 near Moore Haven, FL, along the southwest Lake Okeechobee levee road. Trees within the plots were mechanically uprooted and piled in the center of the plots. One year later

they were burned and the areas harrowed with a disk. Treatments for each plot and numbers of transects and quadrats used in this work are listed in Table 1. After 1 and 2 years, the results of these treatments were assessed.

SUMMARY:

For comparison with melaleuca regeneration in the test plots, trees, as numbers and as stem counts, were determined for an undisturbed site near the test plots. In preparation for harvesting and herbicide treatment, two stands (heads) within Lake Okeechobee were measured and stems counted. To assess melaleuca regeneration after mechanical uprooting of trees and no further site treatment, an area cleared in 1992 was surveyed.

AVAILABILITY OF REPORT:

This report is available on Interlibrary Loan Service from the U.S. Army Engineer Waterways Experiment Station (WES) Library, 3909 Halls Ferry Road, Vicksburg, MS 39180-6199, telephone (601) 634-2355.

To purchase a copy, call the National Technical Information Service (NTIS) at (703) 487-4650. For help in identifying a title for sale, call (703) 487-4780.

NTIS report numbers may also be requested from the WES librarians.

About the Authors:

Dr. Alfred F. Cofrancesco is a research entomologist, WES Environmental Laboratory; Dr. Jean W. Wooten is a botanist, University of Southern Mississippi; and Mr. Harvey L. Jones is a biologist, WES Environmental Laboratory. Point of contact is Dr. Cofrancesco, phone (601) 634-3182.

Contents

Preface	vii
Conversion Factors, Non-SI to SI Units of Measurement	ix
1—Introduction	1
Background	1
Objectives	2
2—Methods	4
Stem Counts in a Melaleuca Stand	6
Stem Counts in Two Melaleuca Stands in Lake Okeechobee	6
Regeneration of Melaleuca After Mechanical Clearing	7
3—Results and Discussion	8
Stem Counts in a Melaleuca Stand	16
Stem Counts in Two Melaleuca Stands in Lake Okeechobee	16
Regeneration of Melaleuca After Mechanical Clearing	17
References	18
Appendix A: Plant Species Present in Sampled Quadrats	A1
SF 298	

List of Tables

Table 1.	Treatment of Melaleuca Test Plots and Numbers of Transects and Quadrats Sampled	3
Table 2.	Estimate Scales for Percent Cover According to Daubenmire (1959) and Dispersion According to Braun Blanquet (1965) ...	5
Table 3.	Totals and Frequency Results for the Species Found in the Study Areas in the Plots	8
Table 4.	Percent Cover Values for the Study Areas in the Plots	9

Table 5.	Species With Greater Than or Equal to 10 Percent of Possible Cover, Growth Form, and Percent of Possible Cover by Plot Number	10
Table 6.	Species With Frequencies Greater Than or Equal to 20 Percent, Relative Percent Frequencies and Percent Possible Cover for the Sampled Areas in the Plots	11
Table 7.	Matrix of Similarity Indices for Presence-Absence (IS_J = Jaccard's Formula) and Percent Cover (IS_E = Ellenberg's Formula) for the Seven Study Areas in the Plots	14
Table 8.	Number of Regenerated Melaleuca Trees in Each Study Area Within the Plots	16
Table A1.	Plant Species Present in Sampled Quadrats	A2

Preface

The work described in this report was authorized by Headquarters, U.S. Army Corps of Engineers (HQUSACE), as part of the Stewardship and Management Task Area of the Wetlands Research Program (WRP). The work was performed under Work Unit 32766, Wetlands Stewardship and Management Demonstration Areas, for which Mr. Chester O. Martin, U.S. Army Engineer Waterways Experiment Station (WES), was the Technical Manager. Ms. Denise White, (CECW-ON) was the WRP Technical Monitor for this work.

Mr. Dave Mathis (CERD-C) was the WRP Coordinator at the Directorate of Research and Development, HQUSACE; Dr. William L. Klesch (CECW-PO) served as the WRP Technical Monitor's Representative; and Dr. Russell F. Theriot, WES, was the Wetlands Program Manager. Mr. Martin was the Task Area Manager.

The work was performed at WES by Dr. Alfred F. Cofrancesco, Principal Investigator, Dr. Jean W. Wooten, University of Southern Mississippi, and Mr. Harvey L. Jones, Aquatic Ecology Branch (AEB), Environmental Laboratory (EL), WES, under the supervision of Dr. Edwin A. Theriot, Chief, AEB; Dr. Conrad J. Kirby, Chief, Ecological Research Division, EL; and Dr. John W. Keeley, Director, EL.

We are appreciative for the assistance of Mr. Ron Medima, USACE, Clewiston, for providing us with information on the site. We also thank Ms. Debbie Miller, Ms. Janis Lanier, Mr. Robert Dew, Jr., Mr. Christopher Bunch, and Mr. Bradley Lewis for assistance with laying out transects, sampling vegetation along the transects, and recording data.

At the time of publication of this report, Director of WES was Dr. Robert W. Whalin. Commander was COL Bruce K. Howard, EN.

This report should be cited as follows:

Cofrancesco, A. F., Jr., Wooten, J. W., and Jones, H. L. (1995).
"Evaluation of mechanical and chemical methods for control of
Melaleuca quinquenervia in southern Florida," Technical Report
WRP-SM-15, U.S. Army Engineer Waterways Experiment Sta-
tion, Vicksburg, MS.

*The contents of this report are not to be used for advertising, publication,
or promotional purposes. Citation of trade names does not constitute an
official endorsement or approval of the use of such commercial products.*

Conversion Factors, Non-SI to SI Units of Measurement

Non-SI units of measurement used in this report can be converted to SI units as follows:

Multiply	By	To Obtain
inches	2.54	centimeters
miles (U.S. statute)	1.609347	kilometers

1 Introduction

Background

Melaleuca quinquenervia (Cav.) Blake, common names melaleuca, cajeput, or punk tree, is native to the Australian region. Trees were first planted in Southern Florida early in the 20th century. Starting in the late 1940s, melaleuca was planted by U.S. Army Engineer personnel at Lake Okeechobee, Florida, to serve as wind barriers and to provide protection for the levees surrounding the lake. The trees have now spread to wetland habitats and were found by a 1979 survey to grow along 88.5 km (53.1 miles) of the 175 km (105 miles) of levee road and to cover approximately 96.8 ha (242 acres) in these perimeter stands (Stocker 1982a) (Figure 1). Additionally, Stocker found that "in the lake proper, melaleuca is found in approximately 420 ha



Figure 1. Mature stand of *Melaleuca quinquenervia* trees at Lake Okeechobee, Florida

(1,050 ac), where it occurs as isolated individuals and as dense colonies (heads) up to 0.6 ha (1.5 ac) in size." This distribution of trees was approximately 0.3 percent of the lake acreage.

It has been estimated that the trees have invaded and occupy 10^6 ha in southern Florida and are spreading to new areas at a rate of about 1,000 ha/year; stands with as many as 5,000 stems per hectare have been reported (DiStefano and Fisher 1983/1984). Attempts to control mature melaleuca at Lake Okeechobee have resulted in reestablishment of dense stands of seedlings (Stocker 1982b). A factor in the aggressive colonizing ability of this species is that trees are often multitemmed and flower up to three times per year. Numerous flowers are borne on the current season's branch growth, the branches continue to grow, and leaves are formed beyond the flowers (Meyers 1983). Approximately 250 very small seeds may be formed in each closed, woody capsule (Woodall 1982). Woodall reported that seed release occurs when the moisture supply to the capsule is interrupted by fire, frost, wind, natural pruning, or human activities. Alexander and Hofstetter (1975) estimated that a single 10-m-tall tree could store over 20 million seeds in its capsules.

Woodall (1983) studied the establishment of melaleuca seedlings in the pine-cypress ecotone of southwest Florida. He found that seeds were long-lived on or in the soil and lost no germination ability after 10 months of shallow burial in a swamp, but seeds buried in a well-drained area lost two-thirds of their viability in this time span. He also reported that burial prevented germination. Moist to saturated soils for several months but rarely flooded provide optimum conditions for tree establishment (Myers 1983).

In order to determine possible mechanisms to control the growth and spread of melaleuca in Lake Okeechobee, the U.S. Army Engineer District, Jacksonville, Florida, sponsored work by Dr. R. K. Stocker of the U.S. Army Engineer Waterways Experiment Station. Five 1982 reports document evaluation of the results of (a) determination of tree distribution at Lake Okeechobee, (b) five herbicides for the control of melaleuca seedlings, (c) four herbicides for the control of mature melaleuca on dredged material islands and levees, (d) five herbicides for control of mature melaleuca by injection, and (e) tree harvesting. These preliminary studies indicated that seedlings and mature trees can be killed by herbicides; injection was about 90-percent effective; and that after 4 months, 66 percent of cut tree stumps resprouted (Stocker 1982a,b,c,d,e).

Objectives

The objectives of this study were to determine the effects of selected treatment methods on vegetation regrowth and to assess their results in control of melaleuca. Seven approximately 3.3-ha plots were established in 1990 near Moore Haven, FL, along the southwest Lake Okeechobee levee road. Trees within the plots were mechanically uprooted and piled in the center of the

plots. One year later they were burned and the areas harrowed with a disk. Treatments for each plot and numbers of transects and quadrats used in this work are listed in Table 1. After 1 and 2 years, the results of these treatments were assessed by methods described below. For comparison with melaleuca regeneration in the test plots, trees, as numbers and as stem counts, were determined for an undisturbed site near the test plots. In preparation for harvesting and herbicide treatment, two stands (heads) within Lake Okeechobee were measured and stems counted. To assess melaleuca regeneration after mechanical uprooting of trees and no further site treatment, an area cleared in 1992 was surveyed.

Table 1 Treatment of Melaleuca Test Plots and Numbers of Transects and Quadrats Sampled				
Plot	Date	Treatment	No. Transects	No. Quadrats
1		Manual clearing of recurring melaleuca	7	56
2	7/15/91	Planted <i>Taxodium</i> and <i>Acer</i> saplings	8	69
3	6/17/91	Rodeo herbicide @ 6.96 L/ha; disk twice on 7/10/91; planted Japanese and Brown Top Millet 2/29/92 @ 56 kg/ha	7	55
4	6/17/91	Rodeo herbicide @ 6.96 L/ha	8	64
5	6/17/91	Control; mechanical clearing	8	64
6	6/17/91	Velpar L herbicide @ 18.8 L/ha	8	63
7	6/17/91	Velpar L herbicide @ 9.38 L/ha	8	65

2 Methods

To eliminate edge effects, 12.8 m were measured at each plot side and plastic pipes were set at the four corners. Surveying flags were placed each 30.5 m from the front corner post to a location where less than 30.5 m existed to the next plot. Survey flags were placed every 15.2 m along each transect line.

At each flagged quadrat point, a 1-m² plastic quadrat was placed with the survey flag in the center (Figure 2). The names of the plant species present were recorded. Percent cover and growth form were estimated using the scales listed in Table 2. All melaleuca in the study areas in each plot were counted, and estimated height and counted numbers of stems noted.




Figure 2. Sampling quadrat at Lake Okeechobee, Florida

Table 2
Estimate Scales for Percent Cover According to Daubenmire
(1959) and Dispersion According to Braun-Blanquet (1965)

Cover			Dispersion	
No.	Percent	Midpoint	No.	Description
1	0-5	2.5	1	Growing solitary
2	5-25	15.0	2	Clumps or dense groups
3	25-50	37.5	3	Small patches or cushions
4	50-75	62.5	4	Small colonies or forming large carpets
5	75-95	85.0	5	Large almost pure population stands
6	95-100	97.5		

The following formulas were used for data analyses from each plot:

- 1) Percent frequency = $\frac{\text{the number of quadrats in which a species occurred}}{\text{the total number of quadrats sampled}} \times 100$
- 2) Relative frequency = $\frac{\text{the number of quadrats in which a species occurred}}{\text{the number of quadrats in which all of the species occurred}} \times 100$
- 3) Percent cover = $\frac{\text{sum of percent cover for the species}}{\text{total m}^2 \text{ sampled in the plot}}$
- 4) Maximum percent possible cover = $\frac{\text{sum of percent cover}}{(\text{maximum possible percent cover (97.5)}) \times \text{no. quadrats sampled in the plot}}$

Each cover scale value was converted to the midpoint percentage of the Daubenmire (1959) class (Table 2).

Similarity between each of the plots was calculated by:

Jaccard's Index of Community Similarity based on presence-absence (Jaccard 1928).

$$5) \quad IS_j = \frac{\text{Number of common species}}{\text{No. unique species in Plot A} + \text{No. unique species in Plot B} + \text{No of common species}} \times 100$$

Ellenberg's Index of Community Similarity based on species quantities (Ellenberg 1956)

$$6) IS_E = \frac{\text{Sum of percent cover of common species Plot A \& B / 2}}{\text{Sum of percent cover of unique Plot A species} + \text{Sum of percent cover of unique Plot B species} + \text{Sum of percent cover of species common to Plot A \& Plot B / 2}} \times 100$$

Each plot was divided into three parts: (a) the area in front of the berm toward the levee, (b) the berm, and (c) the area toward the canal to the rear of the berm. In 1992, soil samples were taken from each of these three areas according to the zigzag technique suggested by the Mississippi Soil Testing Service. For each area in each plot, an aggregate sample was accumulated, sun dried, mixed, and an aliquot removed and sent to the Mississippi Soil Testing Service for analyses of pH, organic matter, phosphorus, potassium, calcium, magnesium, zinc, sulfur, and cation exchange capacity. Results of these analyses were subjected to a one-way analysis of variance procedure.

Stem Counts in a Melaleuca Stand

A population of trees between the levee and rim canal approximately 4.8 km (3 miles) north of the test plots was selected as a "typical undisturbed area." The melaleuca stems within a 0.386-ha (0.952 acre) area were counted in 1992. Trees with stems less than 0.1 m (4 in.) in diameter were recorded as saplings.

Stem Counts in Two Melaleuca Stands in Lake Okeechobee

"Head" 1 was off the airboat trail. Two stakes were affixed to trees distant from the south and west sides of the head. Stake 1 at the south was 230 deg from Stake 2. Stake 1 was 40.5 m (133 ft) and 191 deg from Tree 1 at the southeast corner of the head. Stake 2 was 35 m (115 ft) and 95 deg from Tree 1. The head was measured and the melaleuca stems counted in 1992.

"Head" 2 was northeast of Head 1. Two stakes were affixed to trees distant from the south and east sides of the head. Stake 1 was 128 deg from Stake 2. Stake 1 was 46.9 m (154 ft) and 175 deg from Tree 1 at the southeast corner of the stand. Stake 2 was 39.3 m (129 ft) and 245 deg from Tree 1. The head was measured and the melaleuca stems counted in 1992.

Regeneration of *Melaleuca* After Mechanical Clearing

In July 1992, an approximately 1-mile¹ section between the levee and the rim canal was mechanically cleared of *melaleuca*. The trees were stacked, and no further work was done in the area. In 1993, the numbers of regenerated trees were quantified by establishing 16 transect lines 6 m apart across an area including the space between two piles of trees. These lines were sampled with a square meter quadrat every 6 m.

¹ A table of factors for converting non-SI units of measurement to SI units is presented on page ix.

3 Results and Discussion

A total of 54 transect lines and 436 1-m² quadrats were sampled (Table 1). There were 106 species of plants in the quadrats and 90 species of plants present in each year; 16 were new in 1993, and 16 from 1992 were not present in 1993. The scientific and common names of these species are found in Appendix A. As shown in Table 3, the largest number of species, 53, was in Plot 5, and the smallest, 41, was in Plot 1 during 1992; in 1993, the largest number, 59, was in Plot 4, and the smallest, 39, was in Plot 1.

Table 3
Totals and Frequency Results for the Species Found in the Study Areas in the Plots

Plot No.	No. Species Present		Percent Total Species Present		No. Species with Frequency \geq 20 Percent		Highest Percent Relative Frequency in the Plot	
	1992	1993	1992	1993	1992	1993	1992	1993
1	41	39	44.6	36.1	7	8	9.6	7.9
2	49	52	53.3	48.2	8	8	13.4	10.0
3	48	50	52.2	46.3	10	8	15.0	13.0
4	48	59	52.2	54.6	7	9	17.8	8.1
5	53	51	57.6	47.2	7	9	10.5	7.8
6	45	51	48.9	47.2	7	13	10.4	7.4
7	45	55	48.9	50.9	9	9	11.4	11.3

The plots differed in overall size, elevation, and size of the area used for stacking the trees for burning (the berm). In 1992, the amount of standing water was greatest in Plot 1; all plots were dry during the 1993 study. Plots 6 and 3 had the largest unvegetated areas in 1992, Plots 4 and 5 in 1993; Plots 2 and 4 had 1.2 and 2.5 percent, respectively, less vegetation in 1992 than in 1993; all other plots had increases of from 1.6 (Plot 5) to 44.0 percent (Plot 6). Berm size among the plots varied: Plot 3 was largest; Plot 4 had no berm in transect 8 and was narrower than Plots 1, 2, and 3; smallest in Plots 6 and 7. The berm elevation was conspicuously higher than other areas of the plots. Vegetation on these berms was dominated by *Eupatorium* and species of large, almost pure populations of vines, predominantly *Vigna*.

A cut point for considering the magnitude of times a species was encountered was arbitrarily set at greater than or equal to 20 percent. These numbers of species present per plot were low; five were higher in 1993 and two lower; the number in 1992 was highest at 10 in Plot 3, and in 1993, 13 in Plot 6, indicating that many species were infrequently found in the quadrats (Table 3). In 1992, the highest percent relative frequencies in Plots 1 through 5 were *Eupatorium* and in the last two plots were solitary growing *Baccharis*. In 1993, the highest relative frequencies declined in six of the plots; *Ludwigia octovalvis* was highest in Plots 1 and 2, *Eupatorium* in Plots 3 and 5, *Pluchea* in Plot 4, and *Baccharis* in Plots 6 and 7 (Table 3).

The highest sum of the mean percent cover per square meter of the sampled area per plot was in Plot 1, 87.4 in 1992, 91.8 in 1993; Plot 6 had the lowest in 1992, Plot 5 in 1993 (Table 4). Cover increased in all plots in 1993, most impressively in the herbicide-treated plots, numbers 3, 6, and 7 (Table 4). Few species in any plot had greater than or equal to 75-percent cover (Table 4). Only 8 of the 90 species found in the quadrats had more than 10 percent of possible cover. *Paspalum distichum*, in clumps or dense groups, had over 20 percent of possible cover, which was a reflection of the large frequency and coverage of this species in Plot 1 (Table 5). *Bacopa* had greater cover in more plots in 1993 than in 1992 (Table 5).

Table 4
Percent Cover Values for the Study Areas in the Plots

Plot No.	Sum of Mean Percent Cover/m ² of Sampled Area		No. Species with a Sum of Cover ≥75 Percent	
	1992	1993	1992	1993
1	87.4	91.8	16	14
2	81.3	89.5	19	14
3	57.9	80.2	12	13
4	57.3	63.3	13	11
5	71.1	77.0	17	14
6	50.4	90.0	11	16
7	55.3	81.2	12	18

Species frequencies greater than or equal to 20.0-percent frequency, relative frequencies, and percentages of possible cover are listed in Table 6. These results show that frequency gives little or no indication of cover when determined in quadrates. A species with very small individuals, such as the small *Baccharis* plants in these sampled plots, can produce high frequency values, even though its cover may be low. Likewise, a species with few individuals but large cover over a considerable portion of the sample area will give low frequencies, as was often evident in the data regarding species of *Cyperus* that grew in clumps or dense groups. Of the species listed in Table 6, certain ones repeatedly occur in the plots. These species can be used to characterize each plot either on the basis of frequency or cover.

Table 5
Species With Greater Than or Equal to 10 Percent of Possible
Cover, Growth Form, and Percent of Possible Cover by Plot
Number

Plot No.	Species with ≥ 10 Percent of Possible Cover	Dispersion	Percent of Possible Cover	
			1992	1993
1	<i>Paspalum distichum</i>	5	21.3	25.2
	<i>Ludwigia peruviana</i>	1	10.5	
2	<i>Eupatorium</i>	2	15.4	
	<i>Dactyloctenium</i>	1	10.2	
	<i>Bacopa</i>	4		17.3
	<i>Vigna</i>	5		12.1
3	<i>Vigna</i>	5	11.1	14.9
	<i>Eupatorium</i>	2	10.8	10.0
4	<i>Eupatorium</i>	2	17.6	
	<i>Pluchea</i>	1		11.2
5	<i>Bacopa</i>	4	15.0	12.0
6	<i>Dactyloctenium</i>	1	12.7	
	<i>Bacopa</i>	4		13.1
	<i>Vigna</i>	5		11.4
7	<i>Dactyloctenium</i>	1	10.7	
	<i>Bacopa</i>	4		14.1
	<i>Baccharis</i>	2		12.9
	<i>Vigna</i>	5		11.2
Note: Blanks are for species not meeting the possible cover criteria.				

A matrix of results for calculated similarities of plots based on presence-absence is shown in Table 7. In IS_j , Jaccard's index, equal weight is given to presence and absence of all species. The numerator and denominator change simultaneously. Mueller-Dombois and Ellenberg (1974) state that "experience has shown that there is rarely an IS_j value based on presence which exceeds 50 or 60 percent." Ellenberg's index for percentage cover, IS_E , will be high if the compared plots have common species with high percent coverage. This index reflects not only the number of common and unique species but also the amount of each species present in the plots.

Comparing IS_j values in Table 7, the highest similarities were Plot 5 (Control) and Plot 6 (Velpar at 18.8 L/ha) for 1992, the two herbicide Plots 3 and 7 for 1993. Plots 1 (Manual removal) and 7 (Velpar, 9.38 L/ha) had low similarity indices. Based on presence-absence, the Rodeo herbicide treatments,

Table 6
Species with Frequencies Greater Than or Equal to 20 Percent,
Relative Percent Frequencies, and Percent Possible Cover for
the Sampled Areas in the Plots

Plot No.	Species with Frequencies ≥ 20.0 Percent	Percent Frequency		Relative Percent Frequency		Percent of Possible Cover	
		1992	1993	1992	1993	1992	1993
1	<i>Eupatorium</i>	42.9	30.6	9.6	6.6	5.8	2.4
	<i>Ludwigia octovalvis</i>	42.9	37.1	9.6	7.9	10.5	4.3
	<i>Cyperus surinamensis</i>	32.1		7.3	3.8	0.02	1.1
	<i>Bacopa</i>	25.0	25.8	5.6	5.5	8.2	9.5
	<i>Cyperus polystachos</i>	25.0		5.6	0.3	2.6	0.6
	<i>Dactyloctenium</i>	23.2		5.2	0.0	7.4	0.0
	<i>Paspalum distichum</i>	21.4	33.9	4.8	7.2	21.3	25.2
	<i>Vigna</i>		37.1		7.9		4.3
	<i>Baccharis</i>		30.6		6.6		7.5
	<i>Typha</i>		27.4		5.9		6.2
	<i>Lythrum</i>		21.0		4.5		1.0
2	<i>Eupatorium</i>	63.8	40.3	13.4	7.2	15.4	4.5
	<i>Dactyloctenium</i>	39.1		8.2	0.0	10.2	0.0
	<i>Cyperus polystachos</i>	36.2		7.6	2.6	3.1	0.6
	<i>Ludwigia octovalvis</i>	33.3	56.5	7.0	10.0	6.5	9.9
	<i>Polygonum punctatum</i>	29.0	30.6	6.1	5.4	1.9	1.2
	<i>Baccharis</i>	27.5	30.6	5.8	5.4	2.1	3.8
	<i>Cyperus surinamensis</i>	21.7		4.6	2.3	1.9	0.7
	<i>Bacopa</i>	20.3	38.7	4.3	6.9	3.8	17.7
	<i>Vigna</i>		43.5		7.7		12.0
	<i>Cyperus ligularis</i>		24.2		4.3		9.1
	<i>Juncus megacephalus</i>		21.0		3.7		1.2
3	<i>Eupatorium</i>	87.3	70.5	15.0	13.0	10.8	10.0
	<i>Cyperus polystachos</i>	47.3		8.1	1.2	3.5	0.2
	<i>Baccharis</i>	41.8	49.2	7.2	9.1	3.1	7.5
	<i>Cyperus surinamensis</i>	34.5		5.9	0.6	1.4	0.1
	<i>Ludwigia octovalvis</i>	34.5	42.6	5.9	7.9	6.6	4.0
	<i>Vigna</i>	30.9	45.9	5.3	8.5	11.1	14.9
	<i>Pluchea</i>	30.9		5.3	3.6	0.8	2.6

(Sheet 1 of 3)

Note: Blanks are for species not meeting the frequency criterion.

Table 6 (Continued)

Plot No.	Species with Frequencies ≥ 20.0 Percent	Percent Frequency		Relative Percent Frequency		Percent of Possible Cover	
		1992	1993	1992	1993	1992	1993
3	<i>Dactyloctenium</i>	25.5		4.4	0.0	1.4	0.0
	<i>Eclipta</i>	25.5		4.4	0.3	1.1	0.1
	<i>Panicum bartowense</i>	21.8		3.7	0.0	1.2	0.0
	<i>Bacopa</i>		27.9		5.1		5.7
	<i>Juncus megacephalus</i>		27.9		5.1		2.3
	<i>Cyperus ligularis</i>		23.0		4.2		9.2
	<i>Polygonum punctatum</i>		21.3		3.9		1.0
4	<i>Eupatorium</i>	84.4	37.9	17.8	5.9	17.6	5.6
	<i>Baccharis</i>	43.8	31.0	9.2	4.9	0.9	4.9
	<i>Cyperus polystachos</i>	29.7	20.7	6.3	3.2	2.2	0.5
	<i>Dactyloctenium</i>	29.7		6.3	0.3	1.8	0.1
	<i>Panicum bartowense</i>	26.6		5.6	3.0	8.9	1.9
	<i>Pluchea</i>	21.9	51.7	4.6	8.1	1.6	11.2
	<i>Vigna</i>	21.9	32.8	4.6	5.1	6.5	6.2
	<i>Ludwigia octovalvis</i>		34.5		5.4		3.5
	<i>Phyla</i>		34.5		5.4		3.5
	<i>Bacopa</i>		24.1		3.8		5.1
	<i>Cyperus haspan</i>		24.1		3.8		0.5
5	<i>Eupatorium</i>	48.4	41.7	10.5	7.8	6.1	5.8
	<i>Bacopa</i>	36.0	26.7	7.8	5.0	15.0	11.9
	<i>Cyperus surinamensis</i>	25.0		5.4	2.8	1.2	0.4
	<i>Ambrosia</i>	21.9		4.7	1.6	3.0	0.4
	<i>Baccharis</i>	21.9	41.7	4.7	7.8	0.6	5.0
	<i>Panicum bartowense</i>	21.9	20.0	4.7	3.7	5.4	8.0
	<i>Vigna</i>	21.9	38.3	4.7	7.1	5.0	9.5
	<i>Ludwigia octovalvis</i>		40.0		7.4		3.5
	<i>Mikania</i>		35.0		6.5		3.2
	<i>Lythrum</i>		26.7		5.0		2.0
	<i>Cyperus ligularis</i>		20.0		3.7		4.8
6	<i>Baccharis</i>	50.8	54.0	10.4	7.4	4.0	9.9
	<i>Cyperus polystachos</i>	42.9		8.7	1.9	1.7	0.4
	<i>Dactyloctenium</i>	42.9		8.7	0.0	12.7	0.0

(Sheet 2 of 3)

Table 6 (Concluded)							
Plot No.	Species with Frequencies ≥ 20.0 Percent	Percent Frequency		Relative Percent Frequency		Percent of Possible Cover	
		1992	1993	1992	1993	1992	1993
6	<i>Eupatorium</i>	33.3	31.7	6.8	4.3	0.9	0.8
	<i>Panicum bartowense</i>	28.6		5.8	2.2	8.7	9.0
	<i>Bacopa</i>	23.8	46.0	4.9	6.3	2.0	13.1
	<i>Cyperus surinamensis</i>	20.6		4.2	0.8	0.5	0.4
	<i>Pluchea</i>	20.6	39.6	4.2	5.4	0.7	7.3
	<i>Mikania</i>		34.9		4.8		2.8
	<i>Vigna</i>		33.3		4.6		11.4
	<i>Cynoctonum</i>		30.1		4.1		0.8
	<i>Lythrum</i>		26.9		3.7		0.9
	<i>Ludwigia octovalvis</i>		25.4		3.5		0.9
	<i>Ludwigia microcarpa</i>		23.8		3.3		1.7
	<i>Diodia</i>		22.2		3.0		2.6
	<i>Eleocharis geniculata</i>		20.6		2.8		2.5
	<i>Ipomoea</i>		20.6		2.8		3.6
7	<i>Baccharis</i>	49.2	62.5	11.4	11.3	2.2	12.9
	<i>Ambrosia</i>	30.8		7.1	2.8	7.7	1.0
	<i>Cyperus polystachos</i>	30.8		7.1	2.8	3.1	2.2
	<i>Eupatorium</i>	29.2	28.1	6.8	5.1	0.7	1.3
	<i>Dactyloctenium</i>	29.2		6.8	0.0	10.7	0.0
	<i>Cyperus surinamensis</i>	26.2		6.1	2.3	1.1	0.5
	<i>Pluchea</i>	24.6	23.4	5.7	4.2	0.6	1.3
	<i>Cyperus retrosus</i>	20.0		4.6	0.4	0.5	0.2
	<i>Panicum bartowense</i>	20.0		4.6	1.4	5.4	1.8
	<i>Bacopa</i>		42.1		7.6		14.1
	<i>Mikania</i>		32.8		5.9		3.9
	<i>Vigna</i>		31.2		5.6		11.2
	<i>Ludwigia octovavlis</i>		26.5		4.8		2.1
	<i>Cynoctonum</i>		21.8		4.0		1.0
	<i>Salix</i>		20.3		3.7		4.2
(Sheet 3 of 3)							

Table 7
Matrix of Similarity Indices of Presence-Absence (IS_J = Jaccard's Formula) and Percent Cover (IS_E = Ellenberg's Formula) for the Seven Study Areas In the Plots

Plot No.	Year	Index	Plot Number					
			2	3	4	5	6	7
1	1992	IS_J =	52.4	53.6	53.4	46.9	50.9	43.3
	1993	IS_J =	55.9	55.6	49.3	48.3	57.9	48.4
	1992	IS_E =	84.5	65.8	73.9	72.9	72.9	80.3
	1993	IS_E =	88.9	88.8	80.9	85.9	87.7	83.9
2	1992	IS_J =		51.6	47.0	45.7	44.6	46.9
	1993	IS_J =		62.3	52.1	53.0	52.9	55.9
	1992	IS_E =		82.9	73.6	70.0	74.5	72.4
	1993	IS_E =		90.0	76.2	85.2	91.9	89.3
3	1992	IS_J =			62.7	53.0	55.0	55.0
	1993	IS_J =			54.1	55.6	60.3	65.6
	1992	IS_E =			90.4	75.2	99.0	86.4
	1993	IS_E =			77.3	80.7	81.7	82.6
4	1992	IS_J =				62.9	62.3	52.5
	1993	IS_J =				60.9	62.3	56.2
	1992	IS_E =				82.6	95.0	96.4
	1993	IS_E =				83.2	91.5	88.8
5	1992	IS_J =					63.3	55.6
	1993	IS_J =					60.9	62.1
	1992	IS_E =					85.9	79.6
	1993	IS_E =					89.6	89.5
6	1992	IS_J =						60.7
	1993	IS_J =						64.1
	1992	IS_E =						95.2
	1993	IS_E =						92.9

Plots 3 and 4, were 62.7 similar in 1992, 54.1 in 1993. Disking and planting *Taxodium* and *Acer*, Plot 2, resulted in the lowest similarity with Plot 4, Plot 5, Plot 6, and Plot 7 (Table 7). Of the 21 indices, 6 were less than 50, and all were Plots 1 and 2 comparisons, comprising more than 50 percent of all comparisons made for these two plots in 1992; only three comparisons were less than 50, all Plot 1 in 1993. Sixteen comparisons of 1993 data were higher than those of 1992, indicating that there were more species that occurred more frequently in 1993. Four comparisons for 1993 were less than those of 1992, three involving Plot 4.

Comparisons of plot IS_E percent cover indices in Table 7 indicate that the herbicide treatments alone have high percent cover (88.8 to 95.0) similarities; the greatest 1992 similarity was between Plots 3 and 6, both treated with herbicide. Both these plots had low sum of the mean percent cover per square meter, few species with a sum of cover greater than or equal to 75 (Table 4), and the lowest sum of percent cover indicating that there was much exposed soil. For the 1993 data, Plots 6 and 7 (Velpar) were most similar, 15 values were higher than those in 1992, 6 were less, 3 of them involving Plot 4 or

Plot 3, 3 involving Plot 7. The plots have become more similar in percent cover 2 years after treatment. For 1992, Plots 1 and 2 have the greatest similarity with the highest number of plots in the range of 80 to 85 percent (Plots 3 and 7); for 1992, Plots 1, 2, and 5 (control) had the highest percent cover similarities with the greatest number of plots in the range of 70 to 79.6; for 1993, only two comparisons were in the 70s, Plot 4 with 2 and 3 (Table 7). The lowest 1992 similarity of percent cover was between Plots 1 and 3; for 1993, it was Plots 2 and 4 (Table 7). For 1992, Plot 1 had a high mean percent cover per square meter and Plot 3 a much lower amount (Table 4). Plot 1 had a high number of unique species compared with Plot 3. These differences may reflect the results of differences in treatments or water levels.

The values were used in interpretation of frequency and cover data. Statistical analyses of the soils data were conducted in three ways: (a) aggregating the front, berm, and rear data as single results for each plot, including the control uncleared plot, (b) aggregating the front, berm, and rear data as single results for each plot, excluding the uncleared plot, and (c) aggregating the data from all plots, excluding the uncleared plot that had no berm, into front, berm, and rear. Results indicated that when the soils from an uncleared melaleuca plot were included (method 1 above) in the comparisons, this plot differed from Plots 1 through 7 in its lower pH, higher percent organic matter, and higher sulfur content. This uncleared plot differed from all plots except 5 in its exceptionally high magnesium content (1,941 kg/ha compared with 220s) and from Plot 2 in its high cation exchange capacity (25 meq/100 gm compared with 10). Results from method 2 (above) indicated that Plot 5 differed from Plots 2 and 4 in higher percent organic matter and sulfur content. Results from method 3 (above) indicated that the berm area was much higher in phosphorus and potassium than the front and rear areas of the plots. These high values for the berm areas may partially explain the large amount of vegetation consistently found there.

The trees planted in Plot 2 are living and growing. In wetter areas, they are growing at an impressive rate. The millet planted in Plot 3 did not survive, apparently because of the high water level in the site.

Within Plot 1, *Paspalum distichum* had a high percent frequency and the highest percent of possible cover; this plot had no berm and appeared quite different from the other plots, perhaps because of the high 1992 water level. Plot 2 was distinct because of the manual planting of trees. Plots 3, 4, 6, and 7 showed many similarities probably because of the herbicide treatments. Plot 5 appeared intermediate in its vegetation composition from all other plots.

The numbers of melaleuca trees counted within the study area of each plot are listed in Table 8. The Control, Plots 5 and 2, planted with *Taxodium* and *Acer*, contained the largest numbers of trees. The fewest trees were in Plot 1, the wettest plot in 1992; there were three trees, one at the levee end of a transect and the others on the berm. Velpar at 18.8 L/ha, Plot 6, was the most effective herbicide treatment. These melaleuca regeneration counts suggest that since Plot 5, which has 75 trees, was treated as was Plot 1, which has

Table 8
Number of Regenerated Melaleuca Trees in Each Study Area Within the Plots

Plot Number	Number of Trees
1	3
2	62 (71 stems)
3	46
4	31 (58 stems)
5	75 (82 stems)
6	11
7	28

three trees, perhaps the high water level in Plot 1 prevented regeneration of melaleuca. Uprooting, burning the trees, burying seeds by disking, and not applying herbicide will not control regrowth of melaleuca, as was evidenced by the high number of trees in the control, Plot 5 (Table 8). It is not known if the observed trees are the result of seed germination or sprouting from buried wood.

Stem Counts in a Melaleuca Stand

The approximate size of the area counted was 0.386 ha. The approximate mean counts were 223 stems and 13 saplings (trees with stems less than 0.1 m in diameter). The canopy within this plot was closed; there were few other species growing in the area. These observations indicate that the approximate recorded melaleuca counts may represent a maximum number in many habitats in which this species grows.

Stem Counts in Two Melaleuca Stands in Lake Okeechobee

Head one was approximately 0.102 acre (0.041 ha) and contained approximately 204 stems. There were approximately 146 saplings or 176 stems less than 4 in. in diameter. Head two was approximately 0.134 acres (0.054 ha) and contained approximately 532 stems. There were approximately 204 saplings or 256 stems less than 4 in. in diameter.

Regeneration of *Melaleuca* After Mechanical Clearing

There were a total of 1,238 trees in the 0.0204-ha sampled area. The trees were most frequent near the piles of uprooted trees but extended across a wet ditch toward the levee road. Some measured trees were 3.5 m tall; some trees were in flower.

References

- Alexander, T. R., and Hofstetter, R. H. (1975). "Some current ecological aspects of *Melaleuca quinquenervia* (Cav.) Blake in southern Florida," 39th Annual Meeting of the Florida Academy of Sciences, Lakeland, FL, Abstract.
- Braun-Blanquet, J. (1965). *Plant sociology: The study of plant communities*. C. D. Fuller and H. S. Conard, Transl. rev. and ed., Hafner, London.
- Daubenmire, R. F. (1959). "Canopy coverage method of vegetation analysis," *Northwest Scientist* 33, 43-64.
- DiStefano, J., and Fisher, R. F. (1983/1984). "Invasion potential of *Melaleuca quinquenervia* in southern Florida, U.S.A.," *Forest Ecology and Manage.* 7, 133-141.
- Ellenberg, H. (1956). *Aufgaben und methoden der vegetationskunde*. Eugen Ulmer, Stuttgart.
- Jaccard, P. (1928). "Die statistisch-floristische methode als grundlage der pflanzensoziologie." *Abderhalden, handb. biol. arbeitsmeth.* 11, 165-202.
- Mueller-Dombois, D., and Ellenberg, H. (1974). *Aims and methods of vegetation ecology*. Wiley and Sons, New York.
- Myers, R. L. (1983). "Site susceptibility to invasion by the exotic tree *Melaleuca quinquenervia* in southern Florida," *Jour. Applied Ecology* 20, 645-658.
- Stocker, R. K. (1982a). "Distribution of melaleuca at Lake Okeechobee, Florida," Report to U.S. Army Engineer District, Jacksonville, Jacksonville, FL.
- _____. (1982b). "Herbicidal control of melaleuca seedlings," Report to U.S. Army Engineer District, Jacksonville, Jacksonville, FL.

Stocker, R. K. (1982c). "Herbicidal control of mature *Melaleuca quinquenervia*," Report to U.S. Army Engineer District, Jacksonville, Jacksonville, FL.

_____. (1982d). "Control of *Melaleuca quinquenervia* with injected herbicides," Report to U.S. Army Engineer District, Jacksonville, Jacksonville, FL.

_____. (1982e). "Mechanical harvesting of *Melaleuca quinquenervia* - trial demonstration," Report to U.S. Army Engineer District, Jacksonville, Jacksonville, FL.

Woodall, S. L. (1982). "Seed dispersal in *Melaleuca quinquenervia*," *Florida Scientist* 45, 81-93.

_____. (1983). "Establishment of *Melaleuca quinquenervia* seedlings in the pine-cypress ecotone of southwest Florida," *Florida Scientist* 46, 65-72.

Appendix A

Plant Species Present in Sampled Quadrats

Table A1
Plant Species Present in Sampled Quadrats

Scientific Name	Common Name
<i>Alternanthera philoxeroides</i> (Mart.) Griseb.	Alligator weed
<i>Ambrosia artemisiifolia</i> L.	Common ragweed
<i>Ampelopsis arborea</i> (L.) Rusby	Pepper vine
<i>Andropogon virginicus</i> L.	Broom sedge
<i>Azolla caroliniana</i> Willd.	Mosquito fern
<i>Baccharis halimifolia</i> L.	Eastern baccharis
<i>Bacopa monnieri</i> (L.) Pennel	Water-hyssop
<i>Bidens pilosa</i> L.	Spanishneedles
<i>Canna flaccida</i> Small	Canna
<i>Cardiospermum halicacabrum</i> L.	Ballonvine heartseed
<i>Cenchrus echinatus</i> L.	Sand bur
<i>Centella asiatica</i> (L.) Urban	Spadeleaf
<i>Ceratopteris pteridoides</i> (Hook) Hieron	Floating antler fern
<i>Chamaesyce hyssopifolia</i> (L.) Small	Spurge
<i>Chloris glauca</i> (Chapm.) Wood	Saltmarsh chloris
<i>Chloris petraea</i> Sw.	Tumbleweed windmillgrass
<i>Cicuta mexicana</i> Coult. & Rose	Common waterhemlock
<i>Colocasia esculentum</i> (L.) Schott.	Taro
<i>Commelina diffusa</i> Burm. f.	Spreading dayflower
<i>Conoclinium coelestinum</i> (L.) DC.	Mist flower
<i>Croton glandulosus</i> L.	Croton
<i>Cynoctonum mitreola</i> (L.) Britt.	Lax hompod
<i>Cyperus distinctus</i> Steud.	Flatsedge
<i>Cyperus haspan</i> L.	Sheathed cyperus
<i>Cyperus ligularis</i> L.	Cyperus
<i>Cyperus odoratus</i> L.	Fragrant flatsedge
<i>Cyperus pollardii</i> Britt. ex Small	Sedge
<i>Cyperus polystachos</i> Steud.	Many-spiked flatsedge
<i>Cyperus retrorus</i> Chapm.	Nutsedge
<i>Cyperus surinamensis</i> Rottb.	Tropical flatsedge
<i>Dactyloctenium aegyptium</i> (L.) Richt.	Crowfoot grass
<i>Dichromena colorata</i> (L.) Hitch.	White top
<i>Diodia virginiana</i> L.	Virginia buttonweed
<i>Echinochloa crusgalli</i> (L.) Beauv.	Barnyard grass
(Sheet 1 of 4)	

Table A1 (Continued)	
Scientific name	Common name
<i>Eclipta alba</i> (L.) Hassk.	Common yerbadetajo
<i>Eichhornia crassipes</i> (Mar.) Solms	Waterhyacinth
<i>Eleocharis cellulosa</i> Torr.	Gulfcoast spikerush
<i>Eleocharis geniculata</i> (L.) R. & S.	Spikerush
<i>Emilia fosbergii</i> D. H. Nicholson	Tasselflower
<i>Erechtites hieracifolia</i> (L.) Raf.	Fireweed
<i>Erigeron strigosus</i> Muhl.	Daisy fleabane
<i>Eupatorium capillifolium</i> (Lam.) Small	Dog-fennel
<i>Fimbristylis autumnalis</i> (L.) R. & S.	Slender fimbristylis
<i>Fuirena squarrosa</i> Michx.	Hairy umbrella-sedge
<i>Gallium tinctorum</i> L.	Bye bedstraw
<i>Gaura angustifolia</i> Michx.	Southern gaura
<i>Hydrilla verticillata</i> (L.f.) Royle	Hydrilla
<i>Hydrocotyle umbellata</i> L.	Umbrella pennywort
<i>Ipomoea alba</i> L.	Moon flower
<i>Iresine celosia</i> L.	Iresine
<i>Iris pseudacorus</i> L.	Yellow iris
<i>Juncus biflorus</i> Ell.	Turnflower
<i>Juncus megacephalus</i> M. A. Curtis	Largeheaded rush
<i>Juncus trigonocarpus</i> Steud.	Triangular-fruited rush
<i>Kosteletzkya virginiana</i> (L.) Presl. ex Gray	Seashore-mallow
<i>Lemna</i> sp.	Duckweed
<i>Limnobium spongia</i> (Bosc.) Steud.	American frogbit
<i>Lingernia anagallidea</i> (Michx.) Pennell	Clasping falsepimpernel
<i>Ludwigia linearis</i> Walt.	Narrowleaf seed-box
<i>Ludwigia microcarpa</i> Michx.	Small fruited waterprimrose
<i>Ludwigia octovalvis</i> (Jacq.) Raven	Shrubby waterprimrose
<i>Ludwigia repens</i> Forst.	Marsh purslane
<i>Lythrum alatum</i> var. <i>lanceolatum</i> (Ell.) T. & G.	Winged lythrum
<i>Melaleuca quinquenervia</i> (Cav.) Blake	Punk tree
<i>Mikania scandens</i> (L.) Willd.	Hemp-weed
<i>Momordica charantia</i> L.	Wild balsam apple
<i>Myrica cerifera</i> L.	Wax-myrtle
<i>Panicum bartowense</i> Scribn.	Bartow panicum
(Sheet 2 of 4)	

Table A1 (Continued)	
Scientific name	Common name
<i>Panicum virgatum</i> L.	Switchgrass
<i>Paspalum distichum</i> L.	Knotgrass
<i>Paspalum fimbriatum</i> HBK.	Panama paspalum
<i>Paspalum setaceum</i> Michx.	Thin paspalum
<i>Phragmites australis</i> (Cav.) Trin. ex Steud.	Giant cane
<i>Phyla nodiflora</i> Michx.	Common frog-bit
<i>Physalis viscosa</i> L.	Groundcherry
<i>Pluchea rosea</i> R. K. Godfrey	Purple pluchea
<i>Poinsettia heterophylla</i> (L.) Kl. & Gke.	Poinsettia
<i>Polygala incarnata</i> L.	Milkwort
<i>Polygonum densiflorum</i> Meisn.	Southern smartweed
<i>Polygonum punctatum</i> Ell.	Dotted smartweed
<i>Pontederia cordata</i> L.	Pickereelweed
<i>Portulaca pilosa</i> L.	Purslane
<i>Psilocarya nitens</i> (Vahl.) Wood	Short-beaked bald rush
<i>Ptilimnium capillaceum</i> (Michx.) Raf.	Mock bishopweed
<i>Rhyncholytrum repens</i> (Willd.) C. E. Hubbard	
<i>Rotala ramosior</i> (L.) Koehne	Tooth-cup
<i>Sabatia campanulata</i> (L.) Torr.	Bell shaped sabatia
<i>Sacciolepis striata</i> (L.) Nash	American cupscale
<i>Sagina decumbens</i> (Ell.) T. & G.	Trailing perwort
<i>Sagittaria lancifolia</i> L.	Bulltongue
<i>Salix caroliniana</i> Michx.	Coastal plain willow
<i>Scirpus validus</i> L.	Softstem bullrush
<i>Scoparia dulcis</i> (Ell.) T. & G.	Sweet broomwort
<i>Sesbania</i> sp.	Rattle box
<i>Setaria geniculata</i> (Lam.) Beauv.	Knotroot bristlegrass
<i>Setaria glauca</i> (L.) Beauv.	Yellow foxtail
<i>Sida spinosa</i> L.	Indian mallow
<i>Solanum</i> sp.	Nightshade
<i>Solidago</i> sp.	Goldenrod
<i>Spondias purpurea</i> L.	Hog plum
<i>Sporobolus domingensis</i> (Lam.) Hitch.	Dropseed
<i>Taxodium distichum</i> (L.) Richard	Common baldcypress
(Sheet 3 of 4)	

Table A1 (Concluded)	
Scientific name	Common name
<i>Teucrium canadense</i> L.	Germander
<i>Typha latifolia</i> L.	Common cattail
<i>Verbena scabra</i> Vah.	Verbena
<i>Vigna luteola</i> (Jacq.) Benth.	Yellow cowpea
(Sheet 4 of 4)	

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE December 1995		3. REPORT TYPE AND DATES COVERED Final report
4. TITLE AND SUBTITLE Evaluation of Mechanical and Chemical Methods for Control of <i>Melaleuca quinquenervia</i> in Southern Florida			5. FUNDING NUMBERS	
6. AUTHOR(S) Alfred F. Cofrancesco, Jr., Jean W. Wooten, Harvey L. Jones				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Engineer Waterways Experiment Station 3909 Halls Ferry Road, Vicksburg, MS 39180-6199; University of Southern Mississippi, P.O. Box 5018, Hattiesburg, MS 39401			8. PERFORMING ORGANIZATION REPORT NUMBER Technical Report WRP-SM-15	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Corps of Engineers, Washington, DC 20314-1000			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Available from National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Seven approximately 3.3-ha plots were established near Moore Haven, FL, along the southwest Lake Okeechobee levee road. After mechanical uprooting, stacking, and burning all melaleuca trees, the areas were harrowed with a disk. Two plots were treated with Rodeo herbicide, two with Velpar L, one was planted with common baldcypress (<i>Taxodium distichum</i> (L.) Richard) and red maple (<i>Acer rubrum</i> L.), one was a control with no further treatment, and one had regrowth melaleuca manually removed. One and two years later, results of the these treatments were assessed using transect lines and quadrats. A total of 106 species of plants were found; 90 species each study year were within the quadrats, 16 new in 1993, and 16 absent from the 1992 study. Frequency of species and percent cover increased from 1992 to 1993. Dog-fennel (<i>Eupatorium capillifolium</i> (Lam.)) small and eastern baccharis (<i>Baccharis hamilifolia</i> L.) were cover dominants and most frequent in 1992; these two species were most frequent in 1993 with a variety of species cover dominants. Soil analyses indicated that the control plot differed from the plant and Rodeo herbicide plot in higher percent organic matter and sulfur content. The berm area where the trees were burned was higher in phosphorus and potassium than other areas of the plots. The control plot had the most melaleuca regrowth. The fewest melaleuca trees (three) were counted in the study area of Plot 1, which was not treated with herbicide; it is possible that the 1992 high water level in this plot may have prevented tree regrowth. The highest concentration of Velpar L herbicide treatment plot had 11 trees, the second fewest.				
14. SUBJECT TERMS Cajeput tree Herbicide treatment Mechanical treatment			15. NUMBER OF PAGES 34	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED		18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED		19. SECURITY CLASSIFICATION OF ABSTRACT
20. LIMITATION OF ABSTRACT				

Destroy this report when no longer needed. Do not return it to the originator.